

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁷ : C11D 17/00, 3/00, 1/62, 3/50, 3/20		A1	(11) International Publication Number: WO 00/06690
			(43) International Publication Date: 10 February 2000 (10.02.00)
(21) International Application Number: PCT/US99/17319		(81) Designated States: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HK, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).	
(22) International Filing Date: 29 July 1999 (29.07.99)		Published <i>With international search report.</i>	
(30) Priority Data: 09/126,308 30 July 1998 (30.07.98) US			
(71) Applicant: COLGATE-PALMOLIVE COMPANY [US/US]; 300 Park Avenue, New York, NY 10022 (US).			
(72) Inventors: EWBANK, Eric; Rue Verte 42, B-1950 Kraainem (BE). TUMMERS, Dominique; Rue de Plainevaux 135, B-4100 Seraing (BE).			
(74) Agent: LIEBERMAN, Bernard; Colgate-Palmolive Company, 909 River Road, P.O. Box 1343, Piscataway, NJ 08855-1343 (US).			
(54) Title: WATER-IN-OIL MICROEMULSION FOR PROVIDING COSMETIC ATTRIBUTES TO FABRIC SOFTENING BASE COMPOSITION			
(57) Abstract <p>A perfume-containing water-in-oil microemulsion is provided which is capable of being mixed with an aqueous fabric softener base composition to provide perfume thereto. The use of the defined microemulsion enables the addition of perfume as well as other cosmetic attributes to an aqueous base composition under conditions of low shear while avoiding any problems of gelation or physical instability in the finished fabric softener product.</p> <div style="text-align: right; font-size: 2em;">Y</div>			

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

WATER-IN-OIL MICROEMULSION FOR PROVIDING COSMETIC ATTRIBUTES TO
FABRIC SOFTENING BASE COMPOSITION

Field of Invention

5 This invention relates to a composition and method for "post-adding" cosmetic attributes to a fabric softening base composition at the final stage of its manufacture. More particularly it relates to a water-in-oil microemulsion which is suitable for containing adjuvants such as perfume or colorant and which can be readily mixed with a fabric softening base composition with only a minimum amount of shear and without
10 adversely affecting its physical stability.

Background of the Invention

 Perfumes and colorants are routinely introduced into liquid fabric softening compositions to provide appealing cosmetic attributes for the consumer. The introduction of perfume, for example, to a liquid softening composition, is meant to
15 leave a pleasant and lasting fragrance on the treated fabrics. The colorant is intended to provide a visually pleasing product as well as convey to the consumer an overall impression of softness and quality. From a marketing standpoint, perfumes and colorants are attributes which allow the formulator a degree of flexibility to provide product variety for a given softening technology or for a fixed fabric softening base
20 composition.

 Although perfume, on a weight basis, is invariably a very minor component of a commercial fabric softener composition, its introduction into the composition in an efficient manner is a subject of much concern in the formulation art. This is particularly true for concentrated fabric softeners containing more than about 10% by weight of
25 active softening ingredients insofar as the aqueous softener base compositions tend to gel in the presence of water-immiscible perfume. This tendency of gel formation is

particularly problematic when the water-immiscible perfume is poorly dispersed in the aqueous emulsion which constitutes the softening base composition.

Undesirable thickening or gelation of a softener base composition has significant drawbacks: it reduces the effective shelf life of the final product and adversely affects consumer acceptance for a softener product which upon use has thickened to an unexpectedly viscous mass instead of meeting the consumers expectation of being essentially free-flowing. As a result the prior art has addressed itself to overcoming the pervasive problems of gelation and the formation of high viscosity liquids which accompany the introduction of water-insoluble perfumes into aqueous fabric softener base compositions.

U.S. Patent 5,447,644 to International Flavors & Fragrances (IFF) describes a method to avoid gelation of the softening composition resulting from the introduction of increased levels of perfume into a softener base composition. According to this method there is first formed an aqueous microemulsion by mixing the perfume with a defined nonionic surfactant under conditions of high shear to uniformly disperse the perfume in the surfactant/water mixture. The perfume-containing microemulsion is then mixed with a fabric softener base formulation to form the final fabric softener composition. A high shear mixer is required for this mixing operation.

Accordingly, there remains a need in the art for a method to readily introduce fragrances and colorants into a fabric softener base at the very end of the manufacturing process or even by the consumer himself in the home under very gentle mixing conditions, avoiding the need for high shear mechanical agitation and equipment. Further, there is a need for a method which apart from incorporating perfume or colorants into a base formulation, will do so without adversely impacting the rheological and physical stability of the final product.

Summary of the Invention

In accordance with the present invention there is provided a perfume-containing water-in-oil microemulsion capable of being mixed with a perfume-free fabric softener base composition which is in the form of an aqueous emulsion so as to disperse


5 perfume in said aqueous emulsion under conditions of low shear while avoiding any problem of gelation to thereby provide a physically stable perfume-containing liquid fabric softening composition, said water-in-oil microemulsion comprising:

(a) from about 5% to about 80%, by weight, of a surfactant fabric softener selected from the group consisting of:

10 (i) diester quaternary ammonium compounds having the structural formulae as follows:

(1)

(+)



15 wherein each

A is independently C (O) O - R' or - O (O) C - R';

R is a lower alkyl group having 1 to about 4 carbon atoms;

R' is an alkyl or alkenyl group having 8 to about 22 carbon atoms;

R'' is independently a lower alkyl radical having 1 to about 6 carbon atoms

20 or hydroxyl alkyl group or H;

n is an integer having a value of 1 to about 3; and

X⁻ is a softener compatible anion; and


(2)

+

25 $(R'')_3 - N - (R)_n - (B)_2 \quad X^-$

wherein B is independently A or $(R)_n-A$; and A, R, R" and n are as defined above; and


(3)



5

wherein A, R, R" and n are as defined above; and/or

(ii) diamido ammonium compounds having the formula:



10

wherein n, X⁻ and R' are as defined above, R¹ is a lower alkyl radical having 1 to about 4 carbon atoms or hydrogen, and R is an alkylene radical having 2 to about 4 carbon atoms;

(b) from about 2% to about 50%, by weight, of an organic solvent;

15

(c) from about 5% to about 80%, by weight, of a water-immiscible perfume,

components (a), (b) and (c) comprising the oily phase of said microemulsion;

(d) from 0% to about 1% of a colorant; and

(e) from about 0.5% to about 30%, by weight, of water; said microemulsion

being free of an anionic surfactant, and having a weight ratio of said oily phase to water

20

of from about 3:1 to about 200:1, the percentages of components (a), (b) and (c) and the value of said weight ratio being selected so that the resulting composition forms a water-in-oil microemulsion.

Danielsson and Lindman, "The definition of Microemulsion", Colloids and Surfaces, 3 (1981), 391-392, Elsevier Scientific Publishing Company, have defined a "microemulsion" as 'a system of water, oil and amphiphile which is a single optically isotropic and thermo-dynamically stable liquid solution'. This definition of
5 microemulsion is used herein in describing the present invention.

In accordance with the above definition, a microemulsion has the following characteristics/properties:

- i) Clear
- ii) It is free of lyotropic liquid crystalline system (i.e. it is isotropic)
- 10 iii) It has relatively low viscosity
- iv) It is formed spontaneously by simple mixture/agitation of all components (i.e. it is thermodynamically stable)

More detailed information concerning various aspects of microemulsion systems such as definition, phase behavior, structure, low interfacial tensions and dynamics, is
15 disclosed in R. Zana's article, 'Microemulsions' in Heterogeneous Chemistry Reviews, Vol. 1, 145-157 (1994), edited by John Wiley & Sons Ltd, the disclosure of which is incorporated herein by reference.

The present invention also provides a method for introducing a perfume into a perfume-free fabric softener base composition which is in the form of an aqueous
20 emulsion under conditions of low shear and without adversely affecting its physical stability comprising the steps of:

- (a) providing a perfume-containing water-in-oil microemulsion as defined above; and
- (b) mixing an amount of said water-in-oil microemulsion sufficient to contain
25 the desired amount of perfume with said fabric softener base composition under conditions of low shear thereby causing the inversion of said microemulsion and the

dispersion of the perfume to form a physically stable perfume-containing liquid fabric softening composition.

The water-in-oil microemulsion compositions of the invention are predominantly comprised of the oily phase as defined above, namely, the cationic surfactant fabric softener, the organic solvent and the water-insoluble perfume. To form a composition in the desired phase of a water in oil microemulsion, the amount of water should generally be no greater than about 30%, by weight, and preferably, from about 5% to about 25%, by weight.

The preferred diester quaternary ammonium surfactants for use herein are the dioleoyl diester "quats" represented by equation (1). Particularly preferred is methyl bis-[ethyl(oleoyl)]-2-hydroxyethyl ammonium methyl sulfate, commonly referred to as "DODEQ".

The present invention is predicated on the discovery that cosmetic attributes such as perfume and colorant can be readily "post-added" to an aqueous fabric softening base composition at the final stage of its manufacture by using a water-in-oil microemulsion as herein defined as the vehicle to be added to and mixed with the aqueous emulsion to form the finished product. The water-in-oil microemulsion is incorporated into the softening base composition under conditions of gentle agitation or low shear, using a sufficient amount to provide the desired level of perfume to the finished product. The microemulsion undergoes an inversion upon dilution in the aqueous emulsion which serves to efficiently disperse the perfume or colorant throughout the composition without concomitant problems of gelation or product instability.

Accordingly, the present invention provides important advantages from the standpoint of manufacturing a softening product composition as well as in insuring the integrity of the final product itself. The essence of the present invention is an oil in

water microemulsion which can be readily formed at room temperature with no need for high shear forces or elaborate mixing equipment. And with regard to the finished product, it is clear that the introduction of cosmetic attributes into a softening base composition is accomplished without adversely compromising the rheology or physical stability of the resulting product.

Detailed Description of the Invention

Diester quaternary ammonium surfactant fabric softeners, represented by equation (1) are preferred for use herein and are commercially available from Stepan Co. as Stepanox and from KAO Corp. as Tetranyl but can also be synthesized by the reaction of two moles of a fatty acid with a trialkanolamine, preferably, triethanolamine followed by methylation with dimethyl sulfate or an alkyl halide such as, methyl iodide. In a preferred mode the fatty acid is oleic acid. For economical reasons it has been found that Soya fatty acids are a practical source for this purpose consisting of about 3% myristic acid, about 5% palmitic acid, about 5% palmitoleic acid, 1.5% stearic acid, 72.5% oleic acid and about 13% linoleic acid. Other sources of useful fatty acids are those obtained from the saponification of beef tallow, butter, corn oil, cottonseed oil, lard, olive oil, palm oil, peanut oil, cod liver oil, coconut oil and the like.

A preferred diester quaternary ammonium surfactant fabric softener is methyl bis[ethyl(oleyl)]-2-hydroxyethyl ammonium methyl sulfate. Other diesters useful in the practice of this invention include:

methyl bis-[ethyl(coconut)]-2-hydroxyethyl ammonium methyl sulfate

methyl bis-[ethyl(decyl)]-2-hydroxyethyl ammonium methyl sulfate

methyl bis-[ethyl(dodecyl)]-2-hydroxyethyl ammonium methyl sulfate

methyl bis-[ethyl(lauryl)]-2-hydroxyethyl ammonium methyl sulfate

methyl bis-[ethyl(palmityl)]-2-hydroxyethyl ammonium methyl sulfate

methyl bis-[ethyl(soft-tallow)]-2-hydroxyethyl ammonium methyl sulfate, and the like.

The designation of the terms coconut and soft-tallow indicate mixtures of esters corresponding to the fatty acid source.

In the preparation of the diester quaternary ammonium surfactants, a certain
5 amount of the triester homolog may be produced as an impurity.

The term "perfume" is used herein in its ordinary sense to refer to and include any non water-soluble fragrant substance or mixture of substances including natural (i.e., obtained by extraction of flower, herb, blossom or plant), artificial (i.e., a mixture of natural oils or oil constituents) and synthetic (i.e., a single or mixture of synthetically
10 produced substance) odoriferous substances. Typically perfumes are complex mixtures of blends of various organic compounds, such as, esters, ketones, hydrocarbons, lactones, alcohols, aldehydes, ethers, aromatic compounds and varying amounts of essential oils (e.g., terpenes) such as from about 0% to about 80%, usually from about 10% to 70% by weight, the essential oils themselves being volatile odoriferous
15 compounds and also serving to dissolve the other components of the perfume. The precise composition of the perfume has no particular effect on fabric softening so long as it meets the criteria of water immiscibility and pleasant odor.

Organic solvents suitable for use in this invention include: aliphatic alcohols having 1 to about 6 carbon atoms, such as, ethanol, propanol, isopropanol, n-butanol,
20 isobutanol, t-butanol, n-pentanol, isopentanol, sec-pentanol, n-hexanol, isohexanol, other isomers and the like; aliphatic polyalcohols, such as, ethylene glycol, propylene glycol, butylene glycol, diethylene glycol, dipropylene glycol, 1,4-butanediol, 2-methylpentanediol, hexane triol, tripropylene glycol, pentaerythritol, glycerol, sorbitol, and the like; aliphatic ethers, such as, ethylene glycol monobutyl ether(EGMBE), diethylene
25 glycol monobutyl ether(DEGMBE), diethylene glycol dimethyl ether, triethylene dimethyl ether, ethylene glycol monomethyl ether, propylene glycol monoethyl ether,

dipropylene glycol monomethyl ether, dipropylene glycolpropyl ether(DPnP),
dipropylene glycolbutyl ether(DPnB), tripropylene glycol monomethyl ether, methoxy
methyl butanol, and the like; aliphatic esters, such as, methyl lactate, ethyl lactate,
isopropyl lactate, butyl lactate, dibasic esters of carboxylic acids, ethoxy ethyl acetate,
5 propylene glycol butyl ether acetate, and butoxy ethyl acetate.

Among the preferred organic solvents for use herein are dipropylene glycol
methyl ether (DPM); dipropylene glycol monobutyl ether (DPnB); and ethylene glycol
monobutyl ether (EGMBE).

In the water in oil microemulsion compositions in accordance with the invention,
10 the weight range of cationic surfactant is generally from about 5% to about 80%, and
preferably from about 10% to about 60%, by weight; the weight range of the organic
solvent is generally from about 2% to about 50%, and preferably from about 4% to
about 40%, by weight; and the weight range of the water-immiscible perfume is
generally from about 5% to about 80%, and preferably from about 10% to about 70%,
15 by weight. The water content is generally from about 0.5% to about 30%, and
preferably from about 5% to about 25%, by weight. The weight ratio of the defined oily
phase to water is generally from about 3:1 to about 200:1, preferably from about 5:1 to
50:1, and most preferably from about 5:1 to about 20:1.

The microemulsions of the invention have a particle size between 10 and 100
20 nanometers. At particle sizes below about 50 nanometers, the microemulsions are
generally clear. The microemulsions are formed by simply combining the above-
described components of the composition under gentle agitation or low shear.
Conventional mixing equipment known to those skilled in the art is sufficient for this
purpose. All of the components described herein, both required and optional, must be
25 normally liquid, namely, liquid at ambient room temperatures. Accordingly, there is no
need for heating during the preparation of the microemulsions.

The invention is further described in the examples which follow. All parts and percentages are by weight unless otherwise specified.


Microemulsion compositions/area in a pseudo-ternary phase diagram (i.e. 4 components) were basically determined on a clear/turbid criteria basis by mixing three of the components and titrating with incremental amounts of the fourth component.

Generally, 3.0 g of a mixture consisting of organic solvent (or mixture of), water-immiscible perfume and water, were introduced in a suitable/fixed ratio, and then titrated with DODEQ (dioleoyl diesterquat). Samples were mixed at RT using a magnetic stirrer operating at 100-150 rpm after which they were allowed to stand for few minutes

before visual examination for clarity/turbidity. Clear samples were further examined under a polarizing microscope to check that the liquids were isotropic. By this method, estimated regions for microemulsion compositions were located on pseudo-ternary phase diagrams. Some compositions located inside these microemulsion areas were selected to illustrate the present invention.

Description of chemicals and methods of preparation used in the Examples below.CHEMICALS

The water-in-oil microemulsions contain dioleyl diester quat (DODEQ) as described hereinabove. The formula of DODEQ is as follows:



wherein R₁ and R₂ are C₁₇H₃₄, unsaturated alkyl chains from oleic acid.

DODEQ is commercially available from KAO as Tetranyl or from Stepan Co. as Stepantex. It is synthesized by the reaction of two moles of oleic acid with triethanolamine followed by methylation with dimethyl sulfate.

For purposes of economy, oleic acid may be replaced by olive fatty acids in the synthesis of DODEQ, such olive fatty acids comprising 14% palmitic acid, 2% palmitoleic acid, 2% stearic acid, 64% oleic acid, 16% linoleic acid and 2% linolenic acid. Such material is available from KAO under the name Tetranyl AO-2.

The water-soluble dyes used in the examples are marketed under the names Liquitint Royal Blue, Liquitint Yellow LP and Liquitint Nature Green, by Milliken Chemical company.


The fabric softening base compositions contain two principal softening compounds:

(i) a dialkyl-amidoamine compound commercially available as Rewopal

V3340 from Rewo; and

(ii) a dialkyl esterquat sold as Tetranyl AT175 by KAO.

The dialkyl amidoamine compounds have the general structural formula as follows:



wherein n is an integer of from 1 to 3; R is an alkylene radical having 2 to 4

- 5 carbon atoms; R¹ is an alkyl or alkenyl group having 8 to about 22 carbon atoms; R¹ is a lower alkyl group having 1 to about 4 carbon atoms or hydrogen; and X⁻ is a softener compatible anion.

Method of Preparing Softening Base Composition

- The softening ingredients are each melted, mixed together with stirring and
 10 maintained at about 60 to 70°C. The mixture of molten softening active compounds is added to heated (60-70°C) deionized water with stirring using a 4-pitched blade impeller. The hot water mixture is stirred for 10 to 15 min at about 400 rpm in order to emulsify the molten actives. For concentrated compositions, the stirring speed is increased to 700-800 rpm as the viscosity of the emulsion increases. When required in
 15 the formula composition, hydrochloric acid is first introduced separately in the water prior to the addition of the molten softening ingredients.

- The emulsion is allowed to cool down to 30°C while stirring it at 300-400 rpm. When appropriate, the particle size distribution of the emulsion is further reduced, subjecting the product to high pressure homogenization. This step reduces the
 20 viscosity of the composition. Calcium chloride, preservative, sequestering agent and other optional ingredients such as the thickener, are all introduced sequentially with stirring into the cooled product.

EXAMPLE 1PREPARATION OF PERFUME-CONTAINING MICROEMULSION

Compositions 1 through 6 were prepared following the procedures described above, each composition having varying weight percentages of the four principal components: cationic surfactant; organic solvent; perfume; and water. The ratio of perfume to organic solvent was kept constant at 60:40. The compositions are shown in Table 1 along with observations concerning the appearance of the resulting composition and whether it is within the invention, namely, it formed a water in oil microemulsion, or outside the invention, by forming an emulsion.

TABLE 1

<u>COMPONENT</u>	<u>1</u>	<u>2</u>	<u>3</u>	<u>4</u>	<u>5</u>	<u>6</u>
DODEQ	15%	50%	50%	25%	15%	25%
Organic solvent (DPM) (1)	32	18	14	26	23.6	24
Perfume	48	27	21	39	35.4	36
Water	5	5	15	10	26	15
Appearance of composition	Clear	Clear	Clear	Clear	Turbid	Turbid
Microemulsion (M)/Emulsion (E)	M	M	M	M	E	E

(1) DPM refers to dipropylene glycol methyl ether

As noted in Table 1, Compositions 1-4 are w/o microemulsions within the invention; compositions 5 and 6 are emulsions.

EXAMPLE 2PREPARATION OF W/O MICROEMULSION CONTAINING PERFUMEAND COLORANT

Compositions 7 through 9 were prepared containing a fixed level of perfume and
5 a varying amount of colorant. The compositions are shown below in Table 2 along with
observations concerning the physical appearance of the resulting composition.

TABLE 2

<u>COMPONENT</u>	<u>7</u>	<u>8</u>	<u>9</u>
DODEQ	25%	25%	15%
10 Organic solvent (DPM)	26	26	32
Perfume	39	39	48
Water	9.95	9.5	4.9
Colorant	0.05	0.5	0.1
Appearance of composition	clear(1)	clear(1)	clear(1)
15 Microemulsion (M)/Emulsion (E)	M	M	M

(1) Clear but colored

EXAMPLE 3THE EFFECT OF INCREASING THE LEVEL OF SOLVENT ON THE
MICROEMULSION REGION OF THE PHASE DIAGRAM

20 To demonstrate the effect of increased levels of organic solvent on the region of
microemulsion compositions, compositions 5, 6, 10, 11 and 12 were prepared wherein
the weight ratio of fragrance to organic solvent was varied. The compositions are
shown in Table 3 along with observations concerning the appearance of the resulting
composition.

TABLE 3

<u>COMPONENT</u>	<u>5</u>	<u>10</u>	<u>11</u>	<u>6</u>	<u>12</u>
DODEQ	15%	15%	15%	25%	25%
Perfume + organic solvent (DPM)	59%	59%	59%	60%	60%
5 Ratio of Perfume to organic solvent	60:40	60:50	32:68	60:40	50:50
Water	26%	26%	26%	15%	15%
Appearance of composition	Turbid	Turbid	Clear	Turbid	Clear
Microemulsion (M)/Emulsion (E)	E	E	M	E	M

Compositions 5 and 6 are two compositions from Example 1, which are outside
 10 of the invention but are included in Table 3 as comparative compositions. Table 3
 demonstrates that by changing the ratio of Perfume to organic solvent, it is possible to
 convert a turbid emulsion into a clear fragrance microemulsion suitable for the present
 invention. Thus, a comparison of composition 11, a clear w/o microemulsion, with the
 emulsion compositions 5 and 10 which are outside the invention underscores the
 15 criticality of the perfume to organic solvent ratio. Similarly a comparison of
 compositions 6 and 12 demonstrates that by properly adjusting the ratio of perfume to
 organic solvent, a composition which is outside the invention (composition 6), can be
 reformulated to provide a clear w/o microemulsion (composition 12).

EXAMPLE 4

20 PREPARATION OF W/O MICROEMULSIONS WITH DIFFERENT
CLASSES OF SOLVENTS

Compositions 13 through 18 were prepared to demonstrate the ability to prepare
 microemulsions in accordance with the invention using organic solvents from the
 following classes: ethers, esters, glycols and alkanols. The compositions are shown in
 25 Table 4 below along with observations concerning the appearance of the resulting
 composition.

TABLE 4

<u>COMPONENT</u>	<u>13</u>	<u>14</u>	<u>15</u>	<u>16</u>	<u>17</u>	<u>18</u>
Water	10%	10%	10%	10%	10%	10%
DODEQ	25	25	25	25	25	25
5 Perfume	32.5	32.5	32.5	32.5	32.5	32.5
<u>ORGANIC SOLVENT</u>						
DPM (Ether)	32.5	--	--	--	--	16.25
DPnB (Ether)	--	--	--	--	--	16.25
EGMBE (Ether)	--	32.5	--	--	--	--
10 Hexylene glycol (Glycol)	--	--	32.5	--	--	--
Butanol (Alkanol)	--	--	--	32.5	--	--
Methyl lactate (Ester)	--	--	--	--	32.5	--
<hr/>						
Appearance of Composition	Clear	Clear	Clear	Clear	Clear	Clear
15 Microemulsion (M)/ Emulsion (E)	M	M	M	M	M	M
<hr/>						

EXAMPLE 5DISPERSION OF PERFUME-CONTAINING MICROEMULSION INTO

20

A SOFTENING BASE

The dispersion properties of a perfume-containing microemulsion in accordance with the invention were demonstrated using two different softening base compositions and comparing same with dispersion into 100% water. The perfume micromulsion composition was comprised of the following: 39% perfume; 26% EGMBE organic solvent; 25% DODEQ; 0.48% Liquitint Royal Blue colorant (4% solution); and balance water.

25

A first softening base composition ("AA/EstQ" base) was comprised of the following: 2.74% amidoamine (Rewopal V3340); 1.64% Esterquat (Tetranyl AT1-75); 0.29% glyceryl monooleate; 0.3% hydrochloric acid (25% solution); 0.074% lactic acid (Purac SP80); and balance water.

- 5 The second softening base composition ("EstQ/FA" base) was comprised of the following: 3.9% Esterquat (Tetranyl AT1-75); 0.83% C₁₆₋₁₈ fatty alcohol; 0.2% alcohol ethoxylate (C₁₃₋₁₅ fatty alcohol 20EO); 0.1% amino trimethyl phosphonic acid; 0.063% lactic/lactate buffer solution; 0.20% polyacrylate thickener; and balance water.

- 10 The perfume microemulsion was introduced into each of the above-described softening base compositions at a level of 1.28%, by weight, under very low shear conditions. Mixing was achieved with an Oscell-12 shaker operating for 12 seconds at 700 oscillations per minute. Particle size of the softening base compositions was measured before and after the addition of the perfume microemulsion. For purposes of comparison, dispersion "as is" was measured by post-adding pure perfume to each
- 15 softening base.

TABLE 5PARTICLE SIZE OF SOFTENING BASE BEFORE AND AFTERPOST-ADDITION OF PERFUME

5	<u>SOFTENING BASE</u>	<u>PARTICLE SIZE (μm) BEFORE ADDITION</u>	<u>PARTICLE SIZE (μm) AFTER POST-ADDITION WITH</u>	
			<u>Perfume Microemulsion</u>	<u>Perfume "AS IS"</u>
	AA/EstQ	2.5	2.8	72.0
	EstQ/FA	4.2	5.0	22.0
10	Water	---	24.5	>100

As demonstrated by the data, dispersion of the perfume into the softening base was efficiently carried out using the microemulsion of the invention. The addition of pure perfume ("as is") under the same mixing conditions resulted in an unacceptable dispersion of perfume into the softening base.

15

EXAMPLE 6

The purpose of this Example was to demonstrate the inoperability of a perfume-containing water-in-oil microemulsion which is not formulated in accordance with the invention.

20 The microemulsion composition was comprised of the following:

- (a) 35.8%, by weight, sodium lauryl ether sulfate surfactant;
- (b) 15.4% water;
- (c) 24.4% perfume
- (d) 24.4% DPM organic solvent.

25 The softening base composition was comprised of the following:

- (a) 9.2%, by weight, amidoamine (Rewopal V3340);

- (b) 4.5% Esterquat (Tetranyl AT1-75);
- (c) 1.0% glycerol monooleate;
- (d) 1.4% HCl (25% solution);
- (e) 0.15% calcium chloride (20% solution);
- 5 (f) 0.25% lactic/lactate buffer solution;
- (g) 0.4% Liquitint Royal Blue (4% solution).

Five percent by weight of the perfume-containing w/o microemulsion composition was added to the softening base composition under gentle mixing conditions. A precipitate was observed to form indicating a phase separation and product instability.

I.R. F1376

What is Claimed is:

1. A perfume-containing water-in-oil microemulsion capable of being mixed with a perfume-free fabric softener base composition which is in the form of an aqueous emulsion so as to disperse perfume in said aqueous emulsion under conditions of low shear while avoiding any problem of gelation to thereby provide a physically stable perfume-containing liquid fabric softening composition, said water-in-oil microemulsion comprising:

(a) from about 5% to about 80%, by weight, of a surfactant fabric softener selected from the group consisting of:

(i) diester quaternary ammonium compounds having the structural formulae as follows:

(1)

(+)

15 $(R'')_{4-n} - N - (R-A)_n \quad X^-$

wherein each

A is independently C (O) O - R' or - O (O) C - R';

R is a lower alkyl group having 1 to about 4 carbon atoms;


R' is an alkyl or alkenyl group having 8 to about 22 carbon atoms;

20 R'' is independently a lower alkyl radical having 1 to about 6 carbon atoms or hydroxyl alkyl group or H;

n is an integer having a value of 1 to about 3; and


X⁻ is a softener compatible anion; and

(2)




5 wherein B is independently A or $(R)_n - A$; and A, R, R'' and n are as defined above; and

(3)



wherein A, R, R'' and n are as defined above; and/or

10 (ii) diamido ammonium compounds having the formula:



wherein n, X⁻ and R' are as defined above, R¹ is a lower alkyl radical having 1 to about

15 4 carbon atoms or hydrogen, and R is an alkylene radical having 2 to about 4 carbon atoms;

(b) from about 2% to about 50%, by weight, of an organic solvent;

(c) from about 5% to about 80%, by weight, of a water-immiscible perfume, components (a), (b) and (c) comprising the oily phase of said microemulsion;

20 (d) from 0% to about 1% of a colorant; and

(e) from about 0.5% to about 30%, by weight, of water; said microemulsion being free of an anionic surfactant, and having a weight ratio of said oily phase to water

of from about 3:1 to about 200:1, the percentages of components (a), (b) and (c) and the value of said weight ratio being selected so that the resulting composition forms a water-in-oil microemulsion.

2. A microemulsion as in Claim 1 wherein the diester quaternary ammonium
5 surfactant is methyl bis[ethyl(oleyl)]-2-hydroxyethyl ammonium methyl sulfate.
3. A microemulsion as in Claim 1 wherein said organic solvent is an aliphatic alcohol having from 1 to about 6 carbon atoms.
4. A microemulsion as in Claim 3 wherein said organic solvent is n-butanol.
5. A microemulsion as in Claim 1 wherein said organic solvent is an aliphatic
10 polyalcohol.
6. A microemulsion as in Claim 5 wherein said polyalcohol is hexylene glycol.
7. A microemulsion as in Claim 1 wherein said organic solvent is an aliphatic ether.
8. A microemulsion as in Claim 7 wherein said ether is dipropylene glycol
15 methyl ether.
9. A microemulsion as in Claim 7 wherein said ether is dipropylene glycol n-butyl ether.
10. A microemulsion as in Claim 7 wherein said ether is ethylene glycol
20 monobutyl ether.
11. A microemulsion as in Claim 1 wherein said organic solvent is an aliphatic ester.
12. A microemulsion as in Claim 11 wherein said ester is methyl lactate.
13. A microemulsion as in Claim 1 wherein said fabric softener base
25 composition contains a diester quaternary ammonium softener.

14. A method for introducing a perfume into a perfume-free fabric softener base composition which is in the form of an aqueous emulsion under conditions of low shear and without adversely affecting its physical stability comprising the steps of:

(a) providing a perfume-containing water-in-oil microemulsion as defined in

5 Claim 1; and

(b) mixing an amount of said water-in-oil microemulsion sufficient to contain the desired amount of perfume with said fabric softener base composition under conditions of low shear thereby causing the inversion of said microemulsion and the dispersion of the perfume in said aqueous emulsion to form a physically stable

10 perfume-containing liquid fabric softening composition.

15. A method of imparting softness to fabrics concomitant with a perfume fragrance comprising contacting the fabrics with a softening effective amount of the liquid fabric softening composition formed in Claim 14.

INTERNATIONAL SEARCH REPORT

International Application No
PCT/US 99/17319

A. CLASSIFICATION OF SUBJECT MATTER IPC 7 C11D17/00 C11D3/00 C11D1/62 C11D3/50 C11D3/20		
According to International Patent Classification (IPC) or to both national classification and IPC		
B. FIELDS SEARCHED Minimum documentation searched (classification system followed by classification symbols) IPC 7 C11D		
Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched		
Electronic data base consulted during the international search (name of data base and, where practical, search terms used)		
C. DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	EP 0 813 862 A (HENKEL KGAA) 29 December 1997 (1997-12-29) examples 13,17 ---	1,7, 13-15
A	US 5 525 245 A (GRANDMAIRE JEAN-PAUL ET AL) 11 June 1996 (1996-06-11) claims 1-20; example 29; table 9 ---	1-10,13
A	US 5 447 644 A (GUENIN ERIC P ET AL) 5 September 1995 (1995-09-05) cited in the application claims 1-8 -----	1,14,15
<input type="checkbox"/> Further documents are listed in the continuation of box C. <input checked="" type="checkbox"/> Patent family members are listed in annex.		
* Special categories of cited documents : <div style="display: flex; justify-content: space-between;"> <div> <p>"A" document defining the general state of the art which is not considered to be of particular relevance</p> <p>"E" earlier document but published on or after the international filing date</p> <p>"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)</p> <p>"O" document referring to an oral disclosure, use, exhibition or other means</p> <p>"P" document published prior to the international filing date but later than the priority date claimed</p> </div> <div> <p>"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention</p> <p>"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone</p> <p>"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.</p> <p>"&" document member of the same patent family</p> </div> </div>		
Date of the actual completion of the international search 1 November 1999		Date of mailing of the international search report 25/11/1999
Name and mailing address of the ISA European Patent Office, P.B. 5818 Patentlaan 2 NL - 2280 HV Rijswijk Tel. (+31-70) 340-2040, Tx. 31 651 epo nl, Fax: (+31-70) 340-3016		Authorized officer Loiselet-Taisne, S

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/US 99/17319

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
EP 0813862	A	29-12-1997	DE 19624051 A	18-12-1997
US 5525245	A	11-06-1996	AU 691720 B	21-05-1998
			AU 4525096 A	10-07-1996
			BR 9510472 A	26-05-1998
			CA 2208368 A	27-06-1996
			CN 1173199 A	11-02-1998
			CZ 9701925 A	12-11-1997
			EP 0801672 A	22-10-1997
			FI 972647 A	18-08-1997
			HU 77479 A	28-05-1998
			JP 10511146 T	27-10-1998
			NO 972874 A	13-08-1997
			NZ 300493 A	25-11-1998
			PL 321433 A	08-12-1997
			TR 960592 A	21-07-1996
			WO 9619552 A	27-06-1996
			US 5656585 A	12-08-1997
US 5447644	A	05-09-1995	CA 2148675 A	13-11-1995
			EP 0682104 A	15-11-1995

THIS PAGE BLANK (USPTO)